

MENTAL HEALTH IN NURSING HOMES

SELECTED BIBLIOGRAPHY

Assessment and diagnosis

Cohen-Mansfield J, Deutsch LH. Agitation: subtypes and their mechanisms. *Seminars in Clinical Neuropsychiatry* 1996, 1 (4), 325-339.

Katz IR, Leshner E, Kleban M et al. Clinical features of depression in the nursing home. *International Psychogeriatrics* 1989, 1 (1), 5-15.

Consequences of mental disorder

Lyketsos CG, Steele C, Galik E et al. Physical aggression in dementia patients and its relationship to depression. *American Journal of Psychiatry* 1999, 156 (1), 66-71.

Shinoda-Tagawa T, Leonard R, Pontikas J et al. Resident-to-resident violent incidents in nursing homes. *JAMA* 2004, 291 (5), 591-598.

Suominen K, Henriksson M, Isometsä E. Nursing home suicides: a psychological autopsy study. *International Journal of Geriatric Psychiatry* 2003, 18 (12), 1095-1101.

Cultural and societal issues

Family issues

Gaugler JE, Anderson KA, Zarit SH, Pearlin LI. Family involvement in nursing homes: effects on stress and wellbeing. *Aging and Mental Health* 2004, 8 (1), 65-75.

Hertzberg A, Ekman SL. "We, not them and us?": Views on the relationships between staff and relatives of older people permanently living in nursing homes. *Journal of Advanced Nursing* 2003, 31 (3), 614-622.

Legal and ethical issues

Kirkevold O, Engedal K. Concealment of drugs in food and beverages in nursing homes: cross sectional study. *BMJ* 2005, 330 (7481), 20-22.

Mental health policy

American Geriatrics Society and American Association for Geriatric Psychiatry. Consensus statement on improving the quality of mental health care in US nursing homes: management of depression and behavioral symptoms associated with dementia. *Journal of the American Geriatrics Society* 2003, 51 (9), 1287-1298.

American Geriatrics Society and American Association for Geriatric Psychiatry. The American Geriatrics Society and American Association for Geriatric Psychiatry recommendations for policies in support of quality mental health care in US nursing homes. *Journal of the American Geriatrics Society* 2003, 51 (9), 1299-1304.

Mental health scales

Alexopoulos GS, Abrams RC, Young RC, Shamoian CA. Cornell Scale for Depression in Dementia. *Biological Psychiatry* 1988, 23 (3), 271-284.

Cohen-Mansfield J. Agitated behaviors in the elderly, II: Preliminary results in the cognitively deteriorated. *Journal of the American Geriatrics Society* 1986, 34, 722-7.

Cummings JL, Mega M, Gray K et al. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 1994, 44 (12), 2308-2314.

Reisberg B, Borenstein J, Salob SP et al. Behavioral symptoms in Alzheimer's disease: phenomenology and treatment. *Journal of Clinical Psychiatry* 1987, 48 (suppl 5), 9-15 (Behave-AD).

Yesavage JA, Brink TL, Rose TL et al. Development and validation of a geriatric depression scale: a preliminary report. *Journal of Psychiatric Research* 1983, 17 (1), 37-49.

Models of care and service evaluation

Bartels SJ, Moak GS, Dums AR. Models of mental health services in nursing homes: a review of the literature. *Psychiatric Services* 2002, 53 (11), 1390-1396.

Depla MFIA, Pols J, de Lange J. Integrating mental health care into residential homes for the elderly: an analysis of six Dutch programs for older people with severe and persistent mental illness. *Journal of the American Geriatrics Society* 2003, 51 (9), 1275-1279.

Murphy E. A more ambitious vision for residential long-term care. *International Journal of Geriatric Psychiatry* 1992, 7 (x), 851-852.

Sakauye KM, Camp CJ. Introducing psychiatric care into nursing homes. *Gerontologist* 1992, 32 (6), 849-852.

Schols JMGA, Crebolder HFJM, van Weel C. Nursing home and nursing home physician: the Dutch experience. *Journal of the American Medical Directors Association* 2004, 5 (3), 207-212.

Physical environment and design

Day K, Carreon D, Stump C. The therapeutic design of environments for people with dementia: a review of the empirical research. *Gerontologist* 2000, 40 (4), 398-476.

Zeisel J, Silverstein NM, Hyde J et al. Environmental correlates to behavioral health outcomes in Alzheimer's special care units. *Gerontologist* 2003, 43 (5), 697-711.

Prevalence of mental disorder

Pitkala KH, Laurila JV, Strandberg TE, Tilvis RS. Behavioral symptoms and the administration of psychotropic drugs to aged patients with dementia in nursing homes and in acute geriatric wards. *International Psychogeriatrics* 2004, 16 (1), 61-74.

Smallbrugge M, Jongenelis L, Pot AM, Eefsting JA, Ribbe MW, Beekman ATF. Incidence and outcome of depressive symptoms in nursing home patients in the Netherlands. *American Journal of Geriatric Psychiatry* 2006, 14 (12), 1069-1076.

Teresi J, Abrams R, Holmes D, Ramirez M, Eimicke J. Prevalence of depression and depression recognition in nursing homes. *Social Psychiatry and Psychiatric Epidemiology* 2001, 36 (12), 613-620.

Psychosocial and multi-disciplinary management

Cohen-Mansfield, J. Non-pharmacologic interventions for inappropriate behaviors in dementia: a review, summary, and critique. *American Journal of Geriatric Psychiatry* 2001, 9 (4), 361-381.

Llewellyn-Jones, Baikie KA, Smithers H et al. Multifaceted shared care intervention for late life depression in residential care: randomised controlled trial. *BMJ* 1999, 319, 676-682.

Opie J, Rosewarne R, O'Connor DW. The efficacy of psychosocial approaches to behaviour disorders in dementia: a systematic literature review. *Australian and New Zealand Journal of Psychiatry* 1999, 33 (6), 789-799.

Proctor R, Burns A, Powell HS et al. Behavioural management in nursing and residential homes: a randomised controlled trial. *Lancet* 1999, 354 (9172), 28-29.

Rovner BW, Steele C, Shmueli DS, Folstein MF. A randomized trial of dementia care in nursing homes. *Journal of the American Geriatrics Society* 1996, 44, 7-13.

Psychotropic medication use, efficacy and safety

Draper B, Brodaty H, Low LF et al. Use of psychotropics in Sydney nursing homes: associations with depression, psychosis and behavioral disturbances. *International Psychogeriatrics* 2001, 13 (1), 107-120.

Fossey J, Ballard C, Juszczak E, James I, Alder N, Jacoby R, Howard R. Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial. *BMJ* 2006, 332 (7544), 756-761.

Rabins PV, Lyketsos CG. Antipsychotic drugs in dementia: what should be made of the risks? *JAMA* 2005, 294 (15), 1963-1965.

Sink KM, Holden KF, Yaffe K. Pharmacological treatment of neuropsychiatric symptoms of dementia: a review of the evidence. *JAMA* 2005, 293 (5), 596-608.

Quality of care and quality of life

Ballard C, O'Brien J, James I et al. Quality of life for people with dementia living in residential and nursing home care: the impact of dependency, behaviour and psychotropic drugs. *International Psychogeriatrics* 2001, 13 (1), 93-106.

Samus QM, Rosenblatt A, Steele C et al. The association of neuropsychiatric symptoms and environment with quality of life in assisted living residents with dementia. *Gerontologist* 2005, 45 (1), 19-26.

Staff issues and education

Campbell SL. The concept of attractiveness and older adults with mental illness residing in nursing homes. *Journal of Psychosocial Nursing* 2006, 44 (9), 15-18.

Cole RP, Scott S, Skelton-Robinson M. The effect of challenging behaviour and staff support on the psychological wellbeing of staff working with older adults. *Aging and Mental Health* 2000, 4 (4), 359-65.

Stevens AB, Burgio LD, Bailey E, Burgio K, Paul P, Capilouto E, et al. Teaching and maintaining behaviour management skills with nursing assistants in a nursing home. *Gerontologist* 1998, 38 (3), 379-384.

Textbook

Conn DK, Herrman N, Kaye A, Rewilak D, Schogt B (eds). *Practical Psychiatry in the Long-term Care Home (3rd ed)*. Cambridge, USA: Hogrefe, 2007.